

MYOCARDIAL ISCHEMIA AND INFARCTION

MODERN ERA CORONARY ARTERY BYPASS GRAFT PATENCY: ANGIOGRAPHIC AND FUNCTIONAL OUTCOMES IN THE CASCADE MULTICENTER RANDOMIZED CONTROLLED TRIAL

ACC Oral Contributions

Georgia World Congress Center, Room B404

Tuesday, March 16, 2010, 3:00 p.m.-3:15 p.m.

Session Title: Coronary Artery Bypass Surgery: Contemporary Issues in Graft Survival and Patient Outcomes

Abstract Category: Coronary Artery Bypass Surgery/Innovative Techniques

Presentation Number: 0924-07

Authors: *Alexander Kulik, Pierre Voisine, Patrick Mathieu, Roy G. Masters, Thierry G. Mesana, Michel Le May, Marc Ruel, University of Ottawa Heart Institute, Ottawa, ON, Canada, Hopital Laval, Quebec City, QC, Canada*

Background: In the Clopidogrel after Surgery for Coronary Artery Disease (CASCADE) multi-center randomized controlled trial, patients undergoing primary multi-vessel coronary artery bypass graft surgery (CABG) with at least 2 saphenous vein grafts (SVG) were randomized to receive either aspirin plus clopidogrel or aspirin plus placebo for 1 year postoperatively. The primary outcome was SVG intimal hyperplasia on intravascular ultrasound at 1 year after CABG. The goal of the present substudy was to evaluate graft patency rates and functional outcomes following CABG.

Methods: Ten surgeons at 2 centers performed a mean of 3.5 ± 0.7 grafts (range 2-6 grafts; total 396 grafts) on 113 patients. The mean patient age was 66.5 ± 7.6 years, and 33 patients (29.2%) had diabetes mellitus. One-year clinical follow-up was complete in all patients, and 1-year angiography was completed in 92 patients. In this secondary analysis, we assessed functional outcomes as well as determinants of graft patency 1 year after surgery.

Results: There was no early mortality. One late death occurred from stroke. Four patients required readmission for cardiac ischemic events during the 1 year follow-up. At 1 year, 95.2% of patients were free of angina symptoms. The presence of diabetes did not significantly impact the incidence of cardiac ischemia over the follow-up period (6.1% versus 2.5%, diabetes versus no diabetes, $P=0.58$), and whether patients received more than 3 bypass grafts also did not impact the incidence of cardiac ischemia (4.4% versus 2.2%, ≤ 3 grafts versus >3 grafts, $P=1.0$). Overall 1-year patency was 307 of 322 examined grafts (95.3%). One-year internal thoracic artery graft patency was 98.2%, and 1-year saphenous vein graft patency was 93.8%. The presence of diabetes (96.7% versus 94.8%, diabetes versus no diabetes, $P=0.57$) and the number of grafts performed (94.5% versus 96.2%, ≤ 3 grafts versus >3 grafts, $P=0.60$) did not significantly impact patency.

Conclusions: CABG is associated with excellent angiographic and clinical outcomes in the modern era. The presence of diabetes or more than 3 bypass grafts did not significantly impact graft patency or clinical outcomes in this trial population.